


Sulfonylative Hiyama Cross-Coupling: Development and Mechanistic Insights

A. Adenot,^a J. Char,^a N. von Wolff,^a G. Lefèvre,^a T. Cantat^{a,*}

^a: NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette cedex, France
aurelien.adenot@cea.fr

Due to distinctive structural and electronic features, sulfones have attracted a particular attention over the past few decades, making it a widespread functional group.¹ Present in many contemporary pharmaceuticals and agrochemicals, they are also used as essential intermediates in organic synthesis. Therefore, numerous methodologies have been developed for their preparation.¹ However, the most common methods suffer from significant limitations with harsh reaction conditions or regioselectivity issues. Recently, the insertion of a molecule of sulfur dioxide between two partners was investigated and reactions involving organomagnesium,^{2a} organozinc,^{2b} and organoboron^{2c} compounds were reported. Herein we report a direct single-step palladium-catalyzed synthesis of sulfones involving organosilanes, sulfur dioxide and organohalides. Different mechanistic pathways were envisaged and discussed both from an experimental and theoretical standpoint.



1 N.-W. Liu, S. Liang, G. Manolikakes, *Synthesis*, **2016**, 48, 1939.

2 a) A. S. Deeming, C. J. Russel, A. J. Hennessy, M. C. Willis, *Org. Lett.*, **2014**, 16, 150; b) B. N. Rocke, K. B. Bahnck, M. Herr, S. Lavergne, V. Mascitti, C. Perreault, J. Polivkova, A. Shavnya, *Org. Lett.*, **2014**, 16, 154; c) Y. Chen, M. C. Willis, *Chem. Sci.*, **2017**, 8, 3249.

3 N. von Wolff, J. Char, X. Frogneux, T. Cantat, *Angew. Chem. Int. Ed.*, **2017**, 56, 5616.